



---

Influence of Life Stress on Depression: Moderation by a Polymorphism in the 5-HTT Gene  
Author(s): Avshalom Caspi, Karen Sugden, Terrie E. Moffitt, Alan Taylor, Ian W. Craig, HonaLee Harrington, Joseph McClay, Jonathan Mill, Judy Martin, Antony Braithwaite and Richie Poulton

Reviewed work(s):

Source: *Science*, New Series, Vol. 301, No. 5631 (Jul. 18, 2003), pp. 386-389

Published by: [American Association for the Advancement of Science](#)

Stable URL: <http://www.jstor.org/stable/3834746>

Accessed: 20/09/2012 06:26

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.




*American Association for the Advancement of Science* is collaborating with JSTOR to digitize, preserve and extend access to *Science*.

<http://www.jstor.org>

## REPORTS

**Fig. 4.** PPI of the right ACC. **(A)** All areas are shown that receive a significant context-dependent contribution from right ACC during visuospatial decisions, projected on the same rendered brain as in Fig. 1. Right ACC significantly increased its influence on posterior (28/–72/48,  $t_{\max} = 4.49$ ,  $P < 0.015$ , corrected) and anterior parts of the right IPS (42/–42/44,  $t_{\max} = 4.63$ ,  $P < 0.034$ , corrected) during visuospatial decisions. Note the specificity of this result: Even when the threshold was reduced to  $P < 0.05$ , uncorrected, no other significant clusters were found throughout the brain. **(B)** This schema summarizes the negative findings for right ACC: As indicated by the gray dashed lines, right ACC shows no context-dependent contributions to any left-hemispheric area during visuospatial decisions and none to any left- or right-hemispheric area at all during letter decisions.



tivity (25) that investigated two tasks with right hemisphere dominance demonstrated top-down effects that were specific for the right hemisphere, i.e., from the right middle frontal gyrus (area 46) on right extrastriate areas. It is thus likely that our findings generalize to other lateralized tasks. Although we cannot exclude lateralization contingent on stimulus type in some situations, our results are consistent with previous findings from split-brain patient studies (7) and positron emission tomography (11) showing hemispheric specialization based on task demands. Research on hemispheric specialization should move beyond analyses of asymmetric regional activations and focus more strongly on functional interactions within and between hemispheres.

### References and Notes

1. J. L. Bradshaw, N. C. Nettleton, *Behav. Brain Sci.* **4**, 51 (1981).
2. J. B. Hellige, *Annu. Rev. Psychol.* **41**, 55 (1990).
3. J. C. Marshall, *Behav. Brain Sci.* **4**, 72 (1981).
4. K. Hugdahl, R. J. Davidson, *The Asymmetrical Brain* (MIT Press, Cambridge, MA, 2003).
5. J. Sergent, *Psychol. Bull.* **93**, 481 (1983).
6. C. Chiarello, J. Senehi, M. Soulier, *Neuropsychologia* **4**, 521 (1986).
7. P. M. Corballis, M. G. Funnell, M. S. Gazzaniga, *NeuroReport* **10**, 2183 (1999).
8. C. J. Price, R. J. S. Wise, R. S. J. Frackowiak, *Cereb. Cortex* **6**, 62 (1996).
9. G. R. Fink et al., *Nature* **382**, 626 (1996).
10. L. E. Nyström et al., *Neuroimage* **11**, 424 (2000).
11. S. M. Courtney, L. G. Ungerleider, K. Keil, J. V. Haxby, *Cereb. Cortex* **6**, 39 (1996).
12. J. Levy, C. Trevarthen, *J. Exp. Psychol. Hum. Percept. Perform.* **2**, 299 (1976).
13. M. S. Gazzaniga, *Brain* **123**, 1293 (2000).
14. M. I. Posner, S. E. Petersen, *Annu. Rev. Neurosci.* **13**, 25 (1990).
15. T. Shallice, *From Neuropsychology to Mental Structure* (Cambridge Univ. Press, Cambridge, 1988).
16. R. Desimone, J. Duncan, *Annu. Rev. Neurosci.* **18**, 193 (1995).
17. Materials and methods are available as supporting material on Science Online.
18. Using a  $2 \times 2 \times 2$  factorial design, with letter vs. visuospatial decisions, left vs. right visual fields of presentation, and left vs. right response hand as the three experimental factors, the eight conditions occurred equally often and were varied systematically as blocked conditions in a pseudorandom fashion.

19. A. W. MacDonald, J. D. Cohen, V. A. Stenger, C. S. Carter, *Science* **288**, 1835 (2000).
20. B. J. Casey et al., *Proc. Natl. Acad. Sci. U.S.A.* **97**, 8728 (2000).
21. C. S. Carter et al., *Proc. Natl. Acad. Sci. U.S.A.* **97**, 1944 (2000).

22. G. R. Fink et al., *Brain* **122**, 497 (1999).
23. C. Büchel, K. J. Friston, *Cereb. Cortex* **7**, 768 (1997).
24. A. K. Engel, P. Fries, W. Singer, *Nature Rev. Neurosci.* **2**, 704 (2001).
25. A. R. McIntosh et al., *J. Neurosci.* **14**, 655 (1994).
26. G. Bush, P. Luu, M. I. Posner, *Trends Cogn. Sci.* **4**, 215 (2000).
27. K. J. Friston et al., *NeuroImage* **6**, 218 (1997).
28. C. Cavada, P. S. Goldman-Rakic, *J. Comp. Neurol.* **287**, 393 (1989).
29. D. N. Pandya, G. W. Van Hoesen, M. M. Mesulam, *Exp. Brain Res.* **42**, 319 (1981).
30. This work was supported by the Deutsche Forschungsgemeinschaft (K.Z., G.R.F.), the U.K. Medical Research Council (J.C.M.), and the Wellcome Trust (K.E.S., K.J.F.). Additional support for this Human Brain Project/Neuroinformatics research was provided jointly by the National Institute of Mental Health, National Institute of Neurological Disorders and Stroke, National Institute on Drug Abuse, and the National Cancer Institute. We thank our volunteers, K. Amunts for helpful anatomical discussions, and the radiographers at the Research Center Jülich for technical assistance.

### Supporting Online Material

www.sciencemag.org/cgi/content/full/301/5631/384/DC1  
Materials and Methods  
References

23 April 2003; accepted 6 June 2003

# Influence of Life Stress on Depression: Moderation by a Polymorphism in the 5-HTT Gene

Avshalom Caspi,<sup>1,2</sup> Karen Sugden,<sup>1</sup> Terrie E. Moffitt,<sup>1,2\*</sup>  
Alan Taylor,<sup>1</sup> Ian W. Craig,<sup>1</sup> Honalee Harrington,<sup>2</sup>  
Joseph McClay,<sup>1</sup> Jonathan Mill,<sup>1</sup> Judy Martin,<sup>3</sup>  
Antony Braithwaite,<sup>4</sup> Richie Poulton<sup>3</sup>

In a prospective-longitudinal study of a representative birth cohort, we tested why stressful experiences lead to depression in some people but not in others. A functional polymorphism in the promoter region of the serotonin transporter (5-HTT) gene was found to moderate the influence of stressful life events on depression. Individuals with one or two copies of the short allele of the 5-HTT promoter polymorphism exhibited more depressive symptoms, diagnosable depression, and suicidality in relation to stressful life events than individuals homozygous for the long allele. This epidemiological study thus provides evidence of a gene-by-environment interaction, in which an individual's response to environmental insults is moderated by his or her genetic makeup.

Depression is among the top five leading causes of disability and disease burden throughout the world (1). Across the life span, stressful life events that involve threat, loss, humiliation, or defeat influence the onset and course of depres-

sion (2–5). However, not all people who encounter a stressful life experience succumb to its depressogenic effect. Diathesis-stress theories of depression predict that individuals' sensitivity to stressful events depends on their genetic makeup (6, 7). Behavioral genetics research supports this prediction, documenting that the risk of depression after a stressful event is elevated among people who are at high genetic risk and diminished among those at low genetic risk (8). However, whether specific genes exacerbate or buffer the effect of stressful life events on depression is unknown. In this study, a functional polymorphism in the pro-

<sup>1</sup>Medical Research Council Social, Genetic, and Developmental Psychiatry Research Centre, Institute of Psychiatry, King's College London, PO80 De Crespigny Park, London, SE5 8AF, UK. <sup>2</sup>Department of Psychology, University of Wisconsin, Madison, WI 53706, USA. <sup>3</sup>Dunedin School of Medicine, <sup>4</sup>Department of Pathology, University of Otago, Dunedin, New Zealand.

\*To whom correspondence should be addressed. E-mail: t.moffitt@iop.kcl.ac.uk

motor region of the serotonin transporter gene (*SLC6A4*) was used to characterize genetic vulnerability to depression and to test whether 5-HTT gene variation moderates the influence of life stress on depression.

The serotonin system provides a logical source of candidate genes for depression, because this system is the target of selective serotonin reuptake-inhibitor drugs that are effective in treating depression (9). The serotonin transporter has received particular attention because it is involved in the reuptake of serotonin at brain synapses (10). The promoter activity of the 5-HTT gene, located on 17q11.2, is modified by sequence elements within the proximal 5' regulatory region, designated the 5-HTT gene-linked polymorphic region (5-HTTLPR). The short ("s") allele in the 5-HTTLPR is associated with lower transcriptional efficiency of the promoter compared with the long ("l") allele (11).

Evidence for an association between the short promoter variant and depression is inconclusive (12). Although the 5-HTT gene may not be directly associated with depression, it could moderate the serotonergic response to stress. Three lines of experimental research suggest this hypothesis of a gene-by-environment ( $G \times E$ ) interaction. First, in mice with disrupted 5-HTT, homozygous and heterozygous (5-HTT  $-/-$  and  $+/-$ ) strains exhibited more fearful behavior and greater increases in the stress hormone (plasma) adrenocorticotropin in response to stress compared to homozygous (5-HTT  $+/+$ ) controls, but in the absence of stress no differences related to genotype were observed (13). Second, in rhesus macaques, whose length variation of the 5-HTTLPR is analogous to that of humans, the short allele is associated with decreased serotonergic function [lower cerebrospinal fluid (CSF) 5-hydroxyindoleacetic acid concentrations] among monkeys reared in stressful conditions but not among normally reared monkeys (14). Third, human neuroimaging research suggests that the stress response is mediated by variations in the 5-HTTLPR. Humans with one or two copies of the s allele exhibit greater amygdala neuronal activity to fearful stimuli compared to individuals homozygous for the l allele (15). Taken together, these findings suggest the hypothesis that variations in the 5-HTT gene moderate psychological reactions to stressful experiences.

We tested this  $G \times E$  hypothesis among members of the Dunedin Multidisciplinary Health and Development Study (16). This representative birth cohort of 1037 children (52% male) has been assessed at ages 3, 5, 7, 9, 11, 13, 15, 18, and 21 and was virtually intact (96%) at the age of 26 years. A total of 847 Caucasian non-Maori study members, without stratification confounds, were divided into three groups on the basis of their 5-HTTLPR genotype (11): those with two copies of the s allele (s/s homozygotes;  $n =$

147; 17%), those with one copy of the s allele (s/l heterozygotes;  $n = 435$ ; 51%), and those with two copies of the l allele (l/l homozygotes;  $n = 265$ ; 31%). There was no difference in genotype frequencies between the sexes [ $\chi^2(2) = 0.02$ ,  $P = 0.99$ ]. Stressful life events occurring after the 21st birthday and before the 26th birthday were assessed with the aid of a life-history calendar (17), a highly reliable method for ascertaining life-event histories (18). The 14 events included employment, financial, housing, health, and relationship stressors. Thirty percent of the study members experienced no stressful life events; 25% experienced one event; 20%, two events; 11%, three events; and 15%, four or more events. There were no significant differences between the three genotype groups in the number of life events they experienced,  $F(2,846) = 0.56$ ,  $P = 0.59$ , suggesting that 5-HTTLPR genotype did not influence exposure to stressful life events.

Study members were assessed for past-year depression at age 26 with the use of the Diagnostic Interview Schedule (19), which yields a quantitative measure of depressive symptoms and a categorical diagnosis of a major depressive episode according to *Diagnostic and Statistical Manual of Mental Disorders* (DSM-IV) criteria (20). 17% of study members (58% female versus 42% male; odds ratio = 1.6; 95% confidence interval from 1.1 to 2.2) met criteria for a past-year major depressive episode, which is comparable to age and sex prevalence rates observed in U.S. epidemiological studies (21). In addition, 3% of the study members reported past-year suicide attempts or recurrent thoughts about suicide in the context of a depressive episode. We also collected informant reports about symptoms of depression for 96% of study members at age 26 by mailing a brief questionnaire to persons nominated by each study member as "someone who knows you well."


We used a moderated regression framework (22), with sex as a covariate, to test the association between depression and (i) 5-HTTLPR genotype, (ii) stressful life events, and (iii) their interaction (table S1). The interaction between 5-HTTLPR and life events showed that the effect of life events on self-reports of depression symptoms at age 26 was significantly stronger ( $P = 0.02$ ) among individuals carrying an s allele than among l/l homozygotes (Fig. 1A). We further tested whether life events could predict within-individual increases in depression symptoms over time among individuals with an s allele by statistically controlling for the baseline number of depressive symptoms they had before the life events occurred (table S1). The significant interaction ( $P = 0.05$ ) showed that individuals carrying an s allele whose life events occurred after their 21st birthday experienced increases in depressive symptoms from the age of 21 to 26 years ( $b = 1.55$ ,

$SE = 0.66$ ,  $t = 2.35$ ,  $P = 0.02$  among s/s homozygotes and  $b = 1.25$ ,  $SE = 0.34$ ,  $t = 3.66$ ,  $P < 0.001$  among s/l heterozygotes) whereas l/l homozygotes did not ( $b = 0.17$ ,  $SE = 0.41$ ,  $t = 0.41$ ,  $P = 0.68$ ).

The  $G \times E$  interaction also showed that stressful life events predicted a diagnosis of major depression among carriers of an s allele but not among l/l homozygotes ( $P = 0.056$ , Fig. 1B). We further tested whether life events could predict the onset of new diagnosed depression among carriers of an s allele (table S1). We excluded from analysis study members who were diagnosed with depression before age 21. The significant interaction ( $P = 0.02$ ) showed that life events occurring after their 21st birthdays predicted depression at age 26 among carriers of an s allele who did not have a prior history of depression ( $b = 0.79$ ,  $SE = 0.25$ ,  $z = 3.16$ ,  $P = 0.002$  among s/s homozygotes and  $b = 0.41$ ,  $SE = 0.12$ ,  $z = 3.29$ ,  $P = 0.001$  among s/l heterozygotes) but did not predict onset of new depression among l/l homozygotes ( $b = 0.08$ ,  $SE = 0.20$ ,  $z = 0.42$ ,  $P = 0.67$ ). Further analyses showed that stressful life events predicted suicide ideation or attempt among individuals carrying an s allele but not among l/l homozygotes ( $P = 0.05$ , Fig. 1C). The hypothesized  $G \times E$  interaction was also significant when we predicted informant reports of age-26 depression ( $P < 0.01$ ), an analysis that ruled out the possibility of self-report bias (Fig. 1D). The interaction showed that the effect of life events on informant reports of depression was stronger among individuals carrying an s allele than among l/l homozygotes. These analyses attest that the 5-HTT gene interacts with life events to predict depression symptoms, an increase in symptoms, depression diagnoses, new-onset diagnoses, suicidality, and an informant's report of depressed behavior.

This evidence that 5-HTTLPR variation moderates the effect of life events on depression does not constitute unambiguous evidence of a  $G \times E$  interaction, because exposure to life events may be influenced by genetic factors; if individuals have a heritable tendency to enter situations where they encounter stressful life events, these events may simply be a genetically saturated marker (23, 24). Thus, what we have identified as a gene  $\times$  environment interaction predicting depression could actually reflect a gene  $\times$  "gene" interaction between the 5-HTTLPR and other genes we did not measure. We reasoned that, if our measure of life events represents merely genetic risk, then life events would interact with 5-HTTLPR even if they occurred after the depression episode. However, if our measure of life events represents environmental stress, then the timing of life events relative to depression must follow cause-effect order and life events that

REPORTS




**Fig. 1.** Results of multiple regression analyses estimating the association between number of stressful life events (between ages 21 and 26 years) and depression outcomes at age 26 as a function of 5-HTT genotype. Among the 146 s/s homozygotes, 43 (29%), 37 (25%), 28 (19%), 15 (10%), and 23 (16%) study members experienced zero, one, two, three, and four or more stressful events, respectively. Among the 435 s/l heterozygotes, 141 (32%), 101 (23%), 76 (17%), 49 (11%), and 68 (16%) experienced zero, one, two, three, and four or more stressful events. Among the 264 l/l homozygotes, 79 (29%), 73 (28%), 57 (21%), 26 (10%), and 29 (11%) experienced zero, one, two, three, and four or more stressful events. **(A)** Self-reports of depression symptoms. The main effect of 5-HTTLPR (i.e., an effect not conditional on other variables) was marginally significant ( $b = -0.96$ ,  $SE = 0.52$ ,  $t = 1.86$ ,  $P = 0.06$ ), the main effect of stressful life events was significant ( $b = 1.75$ ,  $SE = 0.23$ ,  $t = 7.45$ ,  $P < 0.001$ ), and the interaction between 5-HTTLPR and life events was in the predicted direction ( $b = -0.89$ ,  $SE = 0.37$ ,  $t = 2.39$ ,  $P = 0.02$ ). The interaction showed that the effect of life events on self-reports of depression symptoms was stronger among individuals carrying an s allele ( $b = 2.52$ ,  $SE = 0.66$ ,  $t = 3.82$ ,  $P < 0.001$  among s/s homozygotes, and  $b = 1.71$ ,  $SE = 0.23$ ,  $t = 5.02$ ,  $P < 0.001$  among s/l heterozygotes) than among l/l homozygotes ( $b = 0.77$ ,  $SE = 0.43$ ,  $t = 1.79$ ,  $P = 0.08$ ). **(B)** Probability of major depressive episode. The main effect of 5-HTTLPR was not significant ( $b = -0.15$ ,  $SE = 0.14$ ,  $z = 1.07$ ,  $P = 0.29$ ), the main effect of life events was significant ( $b = 0.37$ ,  $SE = 0.06$ ,  $z = 5.99$ ,  $P < 0.001$ ), and the  $G \times E$  was in the predicted direction ( $b = -0.19$ ,  $SE = 0.10$ ,  $z = 1.91$ ,  $P = 0.056$ ). Life events predicted a diagnosis of major depression among s carriers ( $b = 0.52$ ,  $SE = 0.16$ ,  $z = 3.28$ ,  $P = 0.001$  among s/s homozygotes, and  $b = 0.39$ ,  $SE = 0.09$ ,  $z = 4.24$ ,  $P < 0.001$  among s/l heterozygotes) but not among l/l homozygotes ( $b = 0.16$ ,  $SE = 0.13$ ,  $z = 1.18$ ,  $P = 0.24$ ). **(C)** Probability of suicide ideation or attempt. The main effect of 5-HTTLPR was not significant ( $b = -0.01$ ,  $SE = 0.28$ ,  $z = 0.01$ ,  $P = 0.99$ ), the main effect of life events was significant ( $b = 0.51$ ,  $SE = 0.13$ ,  $z = 3.96$ ,  $P < 0.001$ ), and the  $G \times E$  interaction was in the predicted direction ( $b = -0.39$ ,  $SE = 0.20$ ,  $t = 1.95$ ,  $P = 0.051$ ). Life events predicted suicide ideation or attempt among s carriers ( $b = 0.48$ ,  $SE = 0.29$ ,  $t = 1.67$ ,  $P = 0.09$  among s/s homozygotes, and  $b = 0.91$ ,  $SE = 0.25$ ,  $z = 3.58$ ,  $P < 0.001$  among s/l heterozygotes) but not among l/l homozygotes ( $b = 0.13$ ,  $SE = 0.26$ ,  $z = 0.49$ ,  $P = 0.62$ ). **(D)** Informant reports of depression. The main effect of 5-HTTLPR was not significant ( $b = -0.06$ ,  $SE = 0.06$ ,  $t = 0.98$ ,  $P = 0.33$ ), the main effect of life events was significant ( $b = 0.23$ ,  $SE = 0.03$ ,  $t = 8.47$ ,  $P < 0.001$ ), and the  $G \times E$  was in the predicted direction ( $b = -0.11$ ,  $SE = 0.04$ ,  $t = 2.54$ ,  $P < 0.01$ ). The effect of life events on depression was stronger among s carriers ( $b = 0.39$ ,  $SE = 0.07$ ,  $t = 5.23$ ,  $P < 0.001$  among s/s homozygotes, and  $b = 0.17$ ,  $SE = 0.04$ ,  $t = 4.51$ ,  $P < 0.001$  among s/l heterozygotes) than among l/l homozygotes ( $b = 0.14$ ,  $SE = 0.05$ ,  $t = 2.69$ ,  $P < 0.01$ ).

occur after depression should not interact with 5-HTTLPR to predict depression. We tested this hypothesis by substituting the age-26 measure of depression with depression assessed in this longitudinal study when study members were 21 and 18 years old, before the occurrence of the measured life events between the ages of 21 and 26 years. Whereas the 5-HTTLPR  $\times$  life events interaction predicted depression at the age of 26 years, this same interaction did not predict

depression reported at age 21 nor at the age of 18 years (table S2), indicating our finding is a true  $G \times E$  interaction.

If 5-HTT genotype moderates the depressogenic influence of stressful life events, it should moderate the effect of life events that occurred not just in adulthood but also of stressful experiences that occurred in earlier developmental periods. Based on this hypothesis, we tested whether adult depression was predicted by the interaction between 5-




**Fig. 2.** Results of regression analysis estimating the association between childhood maltreatment (between the ages 3 and 11 years) and adult depression (ages 18 to 26), as a function of 5-HTT genotype. Among the 147 s/s homozygotes, 92 (63%), 39 (27%), and 16 (11%) study members were in the no maltreatment, probable maltreatment, and severe maltreatment groups, respectively. Among the 435 s/l heterozygotes, 286 (66%), 116 (27%), and 33 (8%) were in the no, probable, and severe maltreatment groups. Among the 265 l/l homozygotes, 172 (65%), 69 (26%), and 24 (9%) were in the no, probable, and severe maltreatment groups. The main effect of 5-HTTLPR was not significant ( $b = -0.14$ ,  $SE = 0.11$ ,  $z = 1.33$ ,  $P = 0.19$ ), the main effect of childhood maltreatment was significant ( $b = 0.30$ ,  $SE = 0.10$ ,  $z = 3.04$ ,  $P = 0.002$ ), and the  $G \times E$  interaction was in the predicted direction ( $b = -0.33$ ,  $SE = 0.16$ ,  $z = 2.01$ ,  $P = 0.05$ ). The interaction showed that childhood stress predicted adult depression only among individuals carrying an s allele ( $b = 0.60$ ,  $SE = 0.26$ ,  $z = 2.31$ ,  $P = 0.02$  among s/s homozygotes, and  $b = 0.45$ ,  $SE = 0.16$ ,  $z = 2.83$ ,  $P = 0.01$  among s/l heterozygotes) and not among l/l homozygotes ( $b = -0.01$ ,  $SE = 0.21$ ,  $z = 0.01$ ,  $P = 0.99$ ).

HTTLPR and childhood maltreatment that occurred during the first decade of life (16, 25). Consistent with the  $G \times E$  hypothesis, the longitudinal prediction from childhood maltreatment to adult depression was significantly moderated by 5-HTTLPR (table S3). The interaction showed ( $P = 0.05$ ) that childhood maltreatment predicted adult depression only among individuals carrying an s allele but not among l/l homozygotes (Fig. 2).

We previously showed that variations in the gene encoding the neurotransmitter-metabolizing enzyme monoamine oxidase A (MAOA) moderate children's sensitivity to maltreatment (25). MAOA has high affinity for 5-HTT, raising the possibility that the protective effect of the l/l allele on psychiatric morbidity is further augmented by the presence of a genotype conferring high MAOA activity (13, 26). However, we found that the moderation of life stress on depression was specific to a polymorphism in the 5-HTT gene, because this effect was observed regardless of the individual's MAOA gene status (tables S4 and S5).

Until this study's findings are replicated, speculation about clinical implications is pre-

**Fig. 3.** The percentage of individuals meeting diagnostic criteria for depression at age 26, as a function of 5-HTT genotype and number of stressful life events between the ages of 21 and 26. The figure shows individuals with either one or two copies of the short allele (left) and individuals



homozygous for the long allele (right). In a hierarchical logistic regression model, the main effect of genotype (coded as s group = 0 and l group = 1) was not significant,  $b = -0.15$ ,  $SE = 0.21$ ,  $z = 0.72$ ,  $P = 0.47$ ; the main effect of number of life events was significant,  $b = 0.34$ ,  $SE = 0.06$ ,  $z = 5.70$ ,  $P < 0.001$ ; and the interaction between genotype and number of life events was significant,  $b = -0.30$ ,  $SE = 0.15$ ,  $z = 1.97$ ,  $P = 0.05$ .

mature. Nonetheless, although carriers of an s 5-HTTLPR allele who experienced four or more life events constituted only 10% of the birth cohort, they accounted for almost one-quarter (23%) of the 133 cases of diagnosed depression. Moreover, among cohort members suffering four or more stressful life events, 33% of individuals with an s allele became depressed, whereas only 17% of the l/l homozygotes developed depression (Fig. 3). Thus, the  $G \times E$ 's attributable risk and predictive sensitivity indicate that more knowledge about the functional properties of the 5-HTT gene may lead to better pharmacological treatments for those already depressed. Although the short 5-HTTLPR variant is too prevalent for discriminatory screening (over half of the Caucasian population has an s allele), a microarray of genes might eventually identify those needing prophylaxis against life's stressful events (27).

Evidence of a direct relation between the 5-HTTLPR and depression has been inconsistent (12), perhaps because prior studies have not considered participants' stress histories. In this study, no direct association between the 5-HTT gene and depression was observed. Previous experimental paradigms, including 5-HTT knockout mice (13), stress-reared rhesus macaques (14), and human functional neuroimaging (15), have shown that the 5-HTT gene can interact with environmental conditions, although these experiments did not address depression. Our study demonstrates that this  $G \times E$  interaction extends to the natural development of depression in a representative sample of humans. However, we could not test hypotheses about brain endophenotypes (28) intermediate between the 5-HTT gene and depression be-

cause of the difficulty of taking CSF or functional magnetic resonance imaging measurements in an epidemiological cohort.

Much genetic research has been guided by the assumption that genes cause diseases, but the expectation that direct paths will be found from gene to disease has not proven fruitful for complex psychiatric disorders (29). Our findings of  $G \times E$  interaction for the 5-HTT gene and another candidate gene, MAOA (25), point to a different, evolutionary model. This model assumes that genetic variants maintained at high prevalence in the population probably act to promote organisms' resistance to environmental pathogens (30). We extend the concept of environmental pathogens to include traumatic, stressful life experiences and propose that the effects of genes may be uncovered when such pathogens are measured (in naturalistic studies) or manipulated (in experimental studies). To date, few linkage studies detect genes, many candidate gene studies fail consistent replication, and genes that replicate account for little variation in the phenotype (29). If replicated, our  $G \times E$  findings will have implications for improving research in psychiatric genetics. Incomplete gene penetrance, a major source of error in linkage pedigrees, can be explained if a gene's effects are expressed only among family members exposed to environmental risk. If risk exposure differs between samples, candidate genes may fail replication. If risk exposure differs among participants within a sample, genes may account for little variation in the phenotype. We speculate that some multifactorial disorders, instead of resulting from variations in many genes of small effect, may result from variations in fewer genes whose effects are conditional on exposure to environmental risks.

## References and Notes

- C. J. Tang, A. D. Lopez, *Lancet* **349**, 1498 (1997).
- G. W. Brown, *Soc. Psychiatry Psychiatr. Epidemiol.* **33**, 363 (1998).
- K. S. Kendler, L. M. Karkowski, C. A. Prescott, *Am. J. Psychiatry* **156**, 837 (1999).
- R. C. Kessler, *Annu. Rev. Psychol.* **48**, 191 (1997).
- D. S. Pine, P. Cohen, J. G. Johnson, J. S. Brook, *J. Affect. Disorders* **68**, 49 (2002).
- E. J. Costello et al., *Biol. Psychiatry* **52**, 529 (2002).
- S. M. Monroe, A. D. Simons, *Psychol. Bull.* **110**, 406 (1991).
- K. S. Kendler et al., *Am. J. Psychiatry* **152**, 833 (1995).
- C. A. Tamminga et al., *Biol. Psychiatry* **52**, 589 (2002).
- K. P. Lesch, M. D. Greenberg, J. D. Higley, A. Bennett, D. L. Murphy, in *Molecular Genetics and the Human Personality*, J. Benjamin, R. P. Ebstein, R. H. Belmaker, Eds. [American Psychiatric Association (APA), Washington, DC, 2002], pp. 109–136.
- K. P. Lesch et al., *Science* **274**, 1527 (1996).
- K. P. Lesch, in *Behavioral Genetics in the Postgenomics Era*, R. Plomin, J. C. DeFries, I. W. Craig, P. McGuffin, Eds. (APA, Washington, DC, 2003), pp. 389–424.
- D. L. Murphy et al., *Brain Res. Bull.* **56**, 487 (2001).
- A. J. Bennett et al., *Mol. Psychiatry* **7**, 188 (2002).
- A. R. Hariri et al., *Science* **297**, 400 (2002).
- Materials and methods are available as supporting material on Science Online.
- A. Caspi et al., *Int. J. Methods Psychiatr. Res.* **6**, 101 (1996).
- R. F. Belli, W. L. Shay, F. P. Stafford, *Public Opin. Q.* **65**, 45 (2001).
- L. N. Robins, L. Cottler, K. Bucholtz, W. Compton, *Diagnostic Interview Schedule for DSM-IV* (Washington University, St. Louis, MO, 1995).
- APA, *Diagnostic and Statistical Manual of Mental Disorders* (APA, Washington, DC, ed. 4, 1994).
- R. C. Kessler, K. A. McGonagle, M. Swartz, D. G. Blazer, C. B. Nelson, *J. Affect. Disorders* **29**, 85 (1993).
- L. S. Aiken, S. G. West, *Multiple Regression: Testing and Interpreting Interactions* (Sage, Thousand Oaks, CA, 1991).
- K. S. Kendler, L. Karkowski-Shuman, *Psychol. Med.* **27**, 539 (1997).
- R. Plomin, C. S. Bergeman, *Behav. Brain Sci.* **14**, 373 (1991).
- A. Caspi et al., *Science* **297**, 851 (2002).
- N. Salichon et al., *J. Neurosci.* **21**, 884 (2001).
- W. E. Evans, M. V. Relling, *Science* **286**, 487 (1999).
- I. I. Gottesman, T. D. Gould, *Am. J. Psychiatry* **160**, 636 (2003).
- D. Hamer, *Science* **298**, 71 (2002).
- A. V. S. Hill, *Br. Med. Bull.* **55**, 401 (1999).
- We thank P. Silva, founder of the Dunedin Multidisciplinary Health and Development Study, Air New Zealand, and the study members, their families, and their friends. Supported by the Health Research Council of New Zealand and the University of Wisconsin Graduate School and by grants from the U.K. Medical Research Council, the William T. Grant Foundation, and the U.S. National Institute of Mental Health (MH49414 and MH45070). T.E.M. is a Royal Society-Wolfson Research Merit Award holder. The study protocol was approved by the institutional review boards of the participating universities.

## Supporting Online Material

www.sciencemag.org/cgi/content/full/301/5631/386/DC1

Materials and Methods  
Tables S1 to S5

27 February 2003; accepted 16 June 2003